

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: Mol Pharmacol 1991 Jun;39(6):711-7

Related Articles, Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov


PubMed Central

Privacy Policy

Phosphodiesterase II, the cGMP-activatable cyclic nucleotide phosphodiesterase, regulates cyclic AMP metabolism in PC12 cells.**Whalin ME, Scammell JG, Strada SJ, Thompson WJ.**

Department of Pharmacology, University of South Alabama College of Medicine, Mobile 36688.

Analysis of cyclic nucleotide phosphodiesterase (PDE) activity in cellular fractions from cultured rat pheochromocytoma (PC12) cells has shown that the predominant hydrolytic activity in both cytosolic and particulate compartments is characteristic of a PDE II, the cGMP-activatable family of PDE isozymes. Cytosolic PDE activity was purified to a high degree utilizing DE-52 anion exchange and cGMP-Sepharose affinity chromatographies. The physicochemical properties of PC12 PDE II were similar to those of PDE II isolated from particulate or soluble fractions of other tissues, including subunit molecular weight of approximately 102,000, activation of cAMP hydrolysis by cGMP, and positive cooperative kinetic behavior for cAMP and cGMP hydrolysis. The potential role of PDE II in regulating cAMP metabolism in intact PC12 cells was studied using an [3H] adenine prelabeling technique. Stimulation of PC12 cell adenosine receptors resulted in a 5-8-fold increase in cAMP accumulation. Removal of the adenosine stimulus by the addition of exogenous adenosine deaminase resulted in a rapid decay of cAMP to prestimulated basal levels within 2 min. Treatment of PC12 cells with atrial natriuretic factor or sodium nitroprusside caused 1) increased intracellular cGMP levels, 2) attenuation of adenosine-stimulated cAMP accumulation, and 3) increased rates of cAMP decay after removal of the adenosine stimulus. Treatment of PC12 cells with HL-725 (a potent inhibitor of isolated PDE II activity in vitro) caused 1) increased basal cAMP accumulation, 2) potentiation of adenosine-stimulated cAMP accumulation, and 3) retardation of the rate of cAMP decay after removal of the adenosine stimulus. HL-725 blocked both the attenuation of cAMP accumulation and the accelerated rate of cAMP decay observed with the cGMP-elevating agents. These results suggest that, in PC12 cells, drugs or hormones that inhibit PDE II or increase intracellular cGMP levels to activate PDE II can modulate cAMP metabolism by altering the catalytic status of the enzyme.



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc

Search for

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Show: Sort Send to

Text Version

☐ 1: Pharmacol Biochem Behav 1999 May;63(1):185-92

[Related Articles, Links](#)

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy


ELSEVIER SCIENCE
FULL-TEXT ARTICLE

Behavioral effects of family-selective inhibitors of cyclic nucleotide phosphodiesterases.

O'Donnell JM, Frith S.

Department of Pharmacology, Louisiana State University School of Medicine, Shreveport 71130, USA.

The effects of family selective inhibitors of phosphodiesterase (PDE1, PDE2, PDE3, PDE4, and PDE5) on the behavior of rats under either a differential-reinforcement-of-low-rate (DRL) 72-s schedule or a variable-interval (VI) 30-s schedule were determined; previous work has shown that antidepressant drugs increase reinforcement rate under long DRL schedules. The PDE4-selective inhibitor rolipram (0.03-0.1 mg/kg) reduced response rate and increased reinforcement rate under the DRL schedule in a dose-dependent manner; similar effects were observed with the tricyclic antidepressant drug desipramine (3-10 mg/kg). Both of these drugs produced biphasic effects on behavior maintained under the VI schedule, increasing response rate at the lower doses tested (rolipram: 0.003 mg/kg; desipramine: 0.03 mg/kg) and decreasing response rate at higher doses (rolipram: 0.1 mg/kg; desipramine: 0.3-18 mg/kg). Of the other PDE inhibitors tested, only the PDE5-selective inhibitor zaprinast (10 mg/kg) produced an antidepressant-like effect on DRL behavior. However, in contrast to the biphasic effects of rolipram and desipramine on VI behavior, zaprinast produced monotonic decreases in response rate (10-30 mg/kg). The PDE2-selective inhibitor trequinsin produced biphasic effects on response rate under the VI schedule, increasing rates at low doses (3-5.6 mg/kg) and decreasing rates at higher doses (18-30 mg/kg). Trequinsin also reduced response rate under the DRL schedule (30 mg/kg); however, the reduction in response rate was not accompanied by increased reinforcement rate. The PDE3-selective inhibitor milrinone (1-10 mg/kg) tended to increase response rates under both schedules while the PDE1-selective inhibitor vinpocetine did not affect behavior at the dose range tested (1-30 mg/kg). These findings suggest that inhibition of PDE4 results in a rather unique pattern of behavioral effects, most notably an antidepressant-like effect on DRL behavior. It remains to be determined if a similar effect produced by zaprinast also implicates PDE5 in the mediation of antidepressant activity or represents an effect of this drug on PDE4 activity.



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc
Search PubMed for [] Go Clear
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Blood Coagul Fibrinolysis 1990 Aug;1(3):247-58

Related Articles, Links

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Arachidonic acid metabolites, ADP and thrombin modulate occlusive thrombus formation over extensive arterial injury in the rat.


De Clerck F, Van Gorp L, Beetens J, Verheyen A, Janssen PA.

Department of Haematology, Janssen Research Foundation, Beerse, Belgium.

Platelet-dependent occlusive thrombosis at sites of deep vessel wall injury elicited by electrical stimulation of rat carotid arteries was significantly reduced by thromboxane A₂ (TXA₂) synthetase inhibition and/or TXA₂/prostaglandin endoperoxide receptor antagonism (ridogrel 1.25 mg/kg i.v.; dazoxiben 5 mg/kg i.v.; sulotroban 20 mg/kg i.v.), by inhibition of ADP-dependent platelet responses (ticlopidine 3 x 200 mg/kg orally) and by anticoagulation (heparin 250 U/kg i.v.; warfarin 1.25 mg/kg i.p.). This points to an involvement of arachidonic acid metabolites, ADP and thrombin as modulators of the thrombotic process. The antithrombotic effect of ridogrel (IC₅₀ = 0.22 mg/kg i.v.) was abolished by cyclooxygenase inhibition (suprofen 5 mg/kg i.v.) but enhanced by cAMP phosphodiesterase inhibition (HL 725 6 micrograms/kg/min i.v.), demonstrating the importance of platelet inhibitory prostanoids such as PGD₂, and prostacyclin formed after TXA₂ synthetase inhibition. High doses of ridogrel (1.25 mg/kg i.v.) producing additional TXA₂/prostaglandin endoperoxide receptor antagonism were more effective than lower doses (0.16 mg/kg i.v.) providing TXA₂ synthetase inhibition alone. The antithrombotic effect of ridogrel, when combined with ticlopidine or heparin, exceeded that of the single compounds, pointing to interactions between arachidonic acid metabolites, ADP and thrombin in the formation of occlusive thrombosis at sites of arterial injury.

PMID: 2151728 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search

PubMed



for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: FEBS Lett 1988 Nov 7;239(2):367-70

Related Articles, Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Nerve growth factor increases the cyclic GMP level and activates the cyclic GMP phosphodiesterase in PC12 cells.**Laasberg T, Pihlak A, Neuman T, Paves H, Saarma M.**

Laboratory of Molecular Genetics, Institute of Chemical Physics and Biophysics, Estonian Academy of Sciences, USSR.

Nerve growth factor (NGF) rapidly increases the cyclic GMP (cGMP) level about 2-3-fold and enhances the cGMP phosphodiesterase (PDE) activity about 2-fold in rat pheochromocytoma PC12 cells. No changes in the level of cyclic AMP (cAMP) and in the activity of cAMP PDE were found. GTP and a nonhydrolysable analog of GTP, GMP-PCP, at 100 microM, were able to mimic the effect of NGF on the cGMP PDE activity. These results suggest that the cGMP system may be one of the second messengers of NGF action in PC12 cells.

PMID: 2460374 [PubMed - indexed for MEDLINE]

Privacy Policy

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Jan 7 2003 16:40:32